

Terms and Conditions

I agree to the condition that this leaflet may be freely reproduced in its entirety by photocopying, print, electronic publishing or any other means provided that **no additions, deletions or any other alterations are made to the text**, and that the text is circulated **for the benefit of patients**.

The Association acknowledges the help of The Royal College of Surgeons of England Patient Liaison Group in the preparation of these leaflets. Patient information produced by the Patient Liaison Group is available at

http://www.rcseng.ac.uk/patient_information/faqs/operation.html/view?searchterm=Patient%20Liaison%20Group

Reaction to ear dressings:

Occasionally the ear may develop an allergic reaction to the dressings in the ear canal. If this happens, the pinna (outer ear) may become swollen and red. You should consult your surgeon so that he can remove the dressing from your ear. The allergic reaction should settle down after a few days.

What happens after the operation?

You will usually go home the day after the operation after the head bandage is removed, or sometimes the same day. There is sometimes some dizziness but this usually settles quickly. The stitches are removed 1 to 2 weeks after the operation.

There may be a small amount of discharge from the ear canal. This usually comes from the ear dressings.

Some of the packing may fall out. If this occurs there is no cause for concern. It is sensible to trim the loose end of packing with scissors and leave the rest in place. The packing in the ear canal will be removed in the ENT Department after 2 or 3 weeks.

You should keep your ear dry. Plug the ear with a cotton wool ball coated with Vaseline when you are having a shower or washing your hair. If the ear becomes more painful or is swollen then you should consult the Ear, Nose and Throat department or your General Practitioner.

You will need to attend the ENT Department occasionally for follow up of your ear for up to 5 years after the operation.

How long will I be off work?

You may need to take 1- 2 weeks off work.

Is there any alternative treatment?

The only way to remove the infection completely is a mastoid operation. In patients who are unfit for surgery, the only alternative is the regular cleaning of the ear by a specialist and the use of antibiotic eardrops. This at best could only reduce the discharge.

**If you have any problems or questions,
please contact:**

*Please insert local department routine and emergency
contact details here*

ABOUT MASTOID SURGERY

ENT.UK is the professional association for Ear, Nose and Throat Surgeons and related professionals in the UK. This information leaflet is to support and not to replace the discussion between you and your specialist. Before you give your consent to the treatment, you should raise any concerns with your specialist.



ENT·UK
The Royal College of Surgeons of England
35-43 Lincoln's Inn Fields
London WC2 3PE
www.entuk.org

Disclaimer

This publication is designed for the information of patients. Whilst every effort has been made to ensure accuracy, the information given may not be comprehensive and patients should not act upon it without seeking professional advice.

Last updated: April 2006
Review due: April 2008
06009

Copyright © 2006 ENT·UK

 **ENT·UK**
BRITISH ASSOCIATION OF OTORHINOLARYNGOLOGISTS
HEAD & NECK SURGEONS

How does the ear work?

The ear consists of the outer, middle and inner ear. The outer ear is covered by skin. The middle ear is covered by a mucus producing membrane. Sound travels through the outer ear and reaches the eardrum, causing it to vibrate. The vibration is transmitted through three tiny bones (ossicles) in the middle ear. The vibration then enters the inner ear where the nerve cells are. The nerve cells within the inner ear are stimulated to produce nerve signals. These nerve signals are carried to the brain, where they are interpreted as sound.

What is the mastoid bone?

The mastoid bone is the bony prominence that can be felt just behind the ear. It contains a number of air spaces, the largest of which is called the antrum. It connects with the air space in the middle ear. Therefore ear diseases in the middle ear can extend into mastoid bone.

Why is mastoid surgery done?

Operations on the mastoid may be necessary when ear infection within the middle ear extends into the mastoid. Most commonly this is a pocket of skin growing from the outer ear into the middle ear, known as cholesteatoma. This causes infection with discharge and some hearing loss. The pocket gets slowly larger, often over a period of many years, and causes gradual erosion of surrounding structures. Erosion of the ossicles can result in hearing loss. The only effective way to get rid of this pocket of skin is surgery.

You may change your mind about the operation at any time, and signing a consent form does not mean that you have to have the operation.

If you would like to have a second opinion about the treatment, you can ask your specialist. He or she

A normal ear drum.



A cholesteatoma.



will not mind arranging this for you. You may wish to ask your own GP to arrange a second opinion with another specialist.

How is mastoid surgery done?

Usually a general anaesthetic is used. There are several ways of doing the operation, depending on the extent of the ear disease and the surgeon. They have various names such as atticotomy and mastoidectomy and take between one and three hours. They involve a cut either above the ear opening or behind the ear.

The bone covering the infection within the mastoid cells is removed. The resultant bony defect is called a mastoid cavity. Some surgeons leave the mastoid cavity open into the ear canal. This allows the surgeons to inspect the mastoid cavity easily. Other surgeons close up the mastoid cavity with bone, cartilage or muscle. You should discuss with your surgeon his/her preferred approach. At the end of the operation, packing will be placed in your ear while it heals.

Does it hurt?

The ear may ache a little but this can be controlled with painkillers provided by the hospital.

How successful is the operation?

The chances of obtaining a dry, trouble free ear from this operation by experienced surgeons are over 80 percent. In some patients it is possible to improve the hearing as well. You should enquire from your surgeon the likelihood of success in your particular case.

Possible complications

There are some risks that you must be aware of before giving consent to this treatment. These potential complications are rare. You should consult your surgeon about the likelihood of problems in your case.

Loss of hearing:

In a small number of patients the hearing may be further impaired due to damage to the inner ear. If the disease has eroded into the inner ear, there may be total loss of hearing in that ear.

Dizziness:

Dizziness is common for a few hours following mastoid surgery and may result in nausea and vomiting. On rare occasions, dizziness is prolonged.

Tinnitus:

Sometimes the patient may notice noise in the ear, in particular if the hearing loss worsens.

Weakness of the face:

The nerve that controls movement of the muscles in the face runs inside the ear and may be damaged during the operation, but this risk is rare. If it happens, the face may lose its movement on one side but it is usually temporary.